arXiv:1303.4964v2 [astro-ph.CO] 23 Apr 2013

The Effects of Temperature Related Viscosity on Cosmological Evolution

Jiaxin Wang [1]* and Xinhe Meng[1,2]†
[1] *Department of Physics, Nankai University, Tianjin 300071, P.R.China and*
[2] *Kavli Institute of Theoretical Physics China,*
CAS, Beijing 100190, China.

(Dated: March 30th 2013)

Bulk viscosity has been intrinsically existing in the observational cosmos evolution with various effects for different cosmological evolution stages endowed with complicated cosmic media. Normally in the idealized "standard cosmology" the physical viscosity effect is often negligent in some extent by assumptions, except for galaxies formation and evolution or the like astro-physics phenomena. Actually we have not fully understood the physical origin and effects of cosmic viscosity, including its practical functions for the universe evolution in reality. In this present article we extend the concept of temperature related viscosity from classical statistical physics to observational cosmology, especially we examine the cosmological effects with possibility of the existence for two kinds of viscosity forms, which are described by the Chapman's relation and Sutherland's formula respectively. With that the unified dark fluid model and also a modification of standard model with viscosity which is named as ΛCDM-V model are constructed physically. While the unified dark fluid model constrained by observational data is closely related with bouncing phenomenon, the ΛCDM-V model is supported more by data-sets fittings with results indicating that temperature related viscosity could not totally explain the rich physics of dark energy phenomena. In addition to the enhancement to cosmic age value, the ΛCDM-V model possesses another two pleasing features: the prediction about the no-rip/singularity future and the mechanism of smooth transition from imperfect cosmological models to perfect ones.

I. INTRODUCTION

The perfect fluid cosmological models are built on the dynamics governed by Einstein's theory of general relativity with the cosmological principle validity at large cosmos scales and the assumption of cosmic compositions modeling as an idealized perfect fluid, which means that all components of the matter-energy in our universe are considered as perfect fluid without any viscosity or which may be interpreted as a result of having taken the gross grain approximation. The well-known ΛCDM model, which is regarded as the most idealized standard description for current cosmology, obtains a constant dark energy term (the famous cosmology constant) that permeates the space everywhere and globally fits the observational data sets very well [29]. But researches have shown that constant dark energy models are not well confirmed by both observations and theoretical considerations [1, 2]. So far the related research work in this line has produced a variety of improved models, and one direction is studying the practical effects of the physically existing viscosity on the cosmic evolutions and its manifestations [3, 12]. It is a natural fact that viscosity exists in the universe evolution history with its eminent role in the thermal plasma as from hot Big Bang like stages and the important effect of turbulence viscosity has been observed both in intra-galactic and inter-cluster medium.

Even though the existence of viscosity is not a problem, what has confused us is its possible origin in the cosmos and how the cosmic viscosity affects the cosmological evolution. In other words, we are searching after physically reasonable explanations and acceptable behaviors of the large-scale existing viscosity in the universe.

It is easy to break down the assumption of perfect fluid by proper modification, but not the cosmological principle which is not only a theoretical principle for large scale cosmology but also an observational feature of our cosmic microwave radiation background. There are two methods for conceiving imperfect fluid models: one obeys the cosmological principle which means the momentum-energy tensor is diagonal in the diagonal metric compatible with the cosmical principle; the other violates the cosmological principle in local region but should be subjected to an average procedure, so that after which the statistical homogeneous and isotropic properties are retrieved at large scales. The first method is widely followed in previous literature, while the second one which is also called back-

---

*Electronic address: jxw@mail.nankai.edu.cn
†Electronic address: xhm@nankai.edu.cn

reaction is famous for its complication and is still in progresses [4].

In this paper, the first method will be adopted in order to build temperature related imperfect cosmological models rather directly. So we arrange the contents as: In the following sub section, before introducing the model buildings we simply review some basic concepts which are closely related with our theme in this present work. In the second section, we will discuss two kinds of imperfect fluid models with a temperature related viscosity origin. The first kind is an unified model where matter and dark energy are indistinguishable clearly or coupled each other, while the second is a modification of the ΛCDM model where the bulk viscosity acts as an extra pressure of total fluid in addition to the normal one. Model constraints with cosmic observational data sets fittings are also included in the second section. We discuss and perform some numerical contrasts for the ΛCDM-V model to ΛCDM model in section three. In the end of this present work, we briefly discuss some features of the new models numerically with the small deviations away from the ΛCDM model and give some conclusions in the fourth section accordingly. For the benefit of the related community we present in details how to do the statistic analysis for the cosmology model buildings with the currently available observational data-sets from direct measuring H(z), SNe Ia from the Union 2.1 and baryon acoustic oscillation (BAO) from large scale surveying as the SDSS (BOSS data-sets) and WMAP-9yr (including the 6dFGS [25] (Beutler *et al.* 2011), Wiggle Z [26] (Blake *et al.*2011), and SDSS III [27] survey results) in the appendix.

A. Einstein's equations

Every homogeneous and isotropic cosmology model is born in the environment of Einstein's dynamic governing equations in the Friedmann Robertson Walker (FRW) metric [31, 34], which read (Since the curvature of the universe is theoretically predicted and observed today to be rather close to zero, so we focus on the flat metric geometry in this article).

$$H^2 = \kappa^2 \rho \tag{1}$$

$$H^2 + \dot{H} = \frac{\ddot{a}}{a} = -\frac{\kappa^2}{2}(\rho + 3p) \tag{2}$$

where $H = \frac{\dot{a}}{a}$ represents the Hubble parameter, alternatively, we usually use $E = H/H_0$ (the dimensionless Hubble parameter) for convenience. Notice that, H_0 is the observed Hubble parameter at present. The combination of the Einstein's component equations (Friedmann equations above) gives the continuous equation

$$\dot{\rho} + 3H(\rho + p) = 0 \tag{3}$$

which means as long as we follow the Einstein's equations, the total energy is automatically conserved. Notice that effective (or total) density ρ and pressure p are concerned in Eqs. (1)–(3).

In addition to the basic functions above, we also pay attention to the expression of equation of state (EOS) parameter which is of vital importance in distinguishing phases of evolution of the universe:

$$w = \frac{p}{\rho} = \gamma(\rho) - 1 \tag{4}$$

In the well-known ΛCDM model, w can be reduced to $-\Omega_\Lambda/E^2$ according to the assumption that radiation occupies so little fraction of the total density that could be neglected in the late stage evolution.

B. viscosity and its classification

Generally speaking, viscosity is still a phenomenological concept generated from the inhomogeneous motion and molecular-like interaction in the fluid [33]. Mathematically the classical viscous stress tensor could be expressed as

$$\sigma_{vis;ij} = \zeta \partial_k v_k \delta_{ij} + \eta(\partial_i v_j + \partial_j v_i - \frac{2}{3}\partial_k v_k \delta_{ij}) \tag{5}$$

where v_k is the k-dimensional component of velocity.

According to the theoretically homogeneous and isotropic property of the large scale observational universe discussed above, we focus on the first term in Eq. (5), namely the bulk viscosity (or the 2nd viscosity) term, while neglecting dissipation and shear viscosity which are incompatible with the cosmological principle. The effective stress tensor of imperfect fluid under cosmological principle now reads

$$\sigma_{ij} = -\delta_{ij}(p - \zeta \partial_k v_k) \tag{6}$$

In the general theory of relativity, the explicit expression of moment-energy tensor then reads

$$T_{\mu\nu} = \rho U_\mu U_\nu + (p - 3H\zeta)h_{\mu\nu} \tag{7}$$

where ρ is energy density, p is isotropic pressure of perfect fluid, $U^\mu = (1, 0, 0, 0)$ is the four-velocity of the cosmic fluid in co-moving coordinates, $h_{\mu\nu} = g_{\mu\nu} + U_\mu U_\nu$ represents the projection tensor, and $3H = \partial_\mu U^\mu$. The similarity could be found between the classically non-relativistic expression (5) and the generally relativistic description about the applications of bulk viscosity in dynamical equations as we presently know.

The characteristic explanation of viscosity could be classified into two types: the dynamical viscosity (bulk and shear viscosity) and the kinematic viscosity which is more usually called turbulence, and is normally represented by a characteristic Reynolds number Re. It is well known that lots of observations support the existence of turbulence, but it is not possible to define a universal Reynolds numbers which characterizes any large-scale complex system. Even though some approximation has been taken in modeling that [12], we should realize that kinematic viscous fluid models could hardly be accepted without considering the back-reaction effects. Meanwhile, the dynamical viscosity is naturally defined to have the feature of unification, which leads to the focus of reaches in the effects of bulk viscosity on cosmological evolution. In this article, we specifically explore the cosmological effects caused by bulk viscosity which is temperature related from molecular-like collision.

C. approaching to viscosity models

The differences between viscous fluid cosmological models are not only due to the different types of viscosity, but also to the different treatments toward the composition of the universe. So far as we know the components highly established both theoretically and observationally of the universe are non-relativistic dust matter (or more specifically, the baryons and dark baryons), cold dark matter mainly, radiation (including relativistic neutrinos and CMB) and the mysterious dark energy. In imperfect fluid models discussed by previous literature researches, either some parts or all of the components of cosmic energy-matter compositions in the universe is assumed having viscous feature. In this present work we take the assumption that viscosity should be a global feature which is more natural and rational as compatible with the cosmological principle, and try to find out whether it is possible to explain dark energy phenomena, which is commonly described by an exotic scalar field such as the quintessence or k-essence, or maybe partly caused by an observational effect of the cosmic media viscosity or due to the gravity modifications to the general theory of relativity at large scales.

According to Eq. (7), the effective pressure which includes the contribution from bulk viscosity reads

$$p = w\rho - 3H\zeta \tag{8}$$

where ζ is the bulk viscosity parameter, and w is the EOS parameter of cosmic fluid except for the viscosity. For unified fluid model (we neglect radiation since it contributes little to the late-stage evolution scenario), it is assumed that the universe only consists of pressless dust (baryons plus cold dark matter). The EOS parameter w in Eq. (8) should consequently be zero, and then the cosmological expansion is caused mainly by the negative pressure $-3H\zeta$ provided by viscosity only.

Much efforts have been made to study imperfect models with simply constant viscosity parameter in previous studies. One drawback of that assumption is that it has overemphasized the differences between perfect fluid models and imperfect ones. We believe that the perfect fluid universe should be an evolutional approximation of imperfect fluid universe when it goes infinities, and the evolutional behavior of viscosity should be functioning.

In this article we try to extend the concept which is one possible origin of cosmic viscosity—molecular-like (galaxies and galaxies collision, for example) interaction—to cosmology researches. On large scales we assume, that most of the visible cosmic matter is effectively in a thermal equilibrium state characterized by a certain temperature, and the interaction between galaxies or matter clumps could be approximately described in a statistical way.

For imperfect fluid, whose viscosity is generated from molecular-like interaction and characterized by macroscopic temperature of the environment, the bulk viscosity can be expressed by a function $\zeta(T)$ as we have just discussed above. The CMB spectrum indicates that the universe has reached a thermal equilibrium state with a typical temperature, which means that the collision or interaction in the equilibrium state can be represented by the average temperature of ambient photons which pass through the universe and frequently interact with all luminous matter.

Under those considerations, we can put forward in another way that global viscosity is naturally introduced by the gross grain approximation and related to the CMB temperature at various stages (which will be discussed briefly in the next subsection).

In this paper, we adopt two appropriate molecular-like collision/ interaction models for cosmology study, from which the viscosity is physically generated; one is based on the Chapman-Enskog formula [32] for dilute fluid which reads

$$\zeta = 2.67 \times 10^{-6} \frac{(mT)^{\frac{1}{2}}}{\sigma^2 \omega(T^*)} \tag{9}$$

where σ is collision diameter, ω is collision integral, $T^* = kT/\epsilon$ with the maximum energy of attraction ϵ divided by Boltzmann constant k, and m is molecular like mass.

The other one is the Sutherland's formula [32] which reads

$$\zeta = \zeta'(\frac{T}{T'})^{\frac{3}{2}} \frac{T' + S}{T + S} \tag{10}$$

where S is Sutherland's constant, the prime represents reference values here.

The above two models are both semi-theoretical or empirical and with first order approximations. The Chapman-Enskog relation assumes that molecules possess only translational kinetic energy, while Sutherland's formula assumes molecules are smooth rigid elastic spheres surrounded by fields of attractive force. Both forms are available for building cosmological models under proper simplification, since the empirically determined parameters are impossible to be given precisely on cosmic large scales. At low temperature ($T \ll 300K$), a general equation which can approximately represent the main feature of Chapman's and Sutherland's formulae reads

$$\zeta = \zeta_0 T^{\alpha} \tag{11}$$

where ζ_0 is considered as a simplified coefficient, and α equals $\frac{1}{2}$ or $\frac{3}{2}$ which are chosen specifically in order to physically represent Chapman's or Sutherland's limit formula respectively. Theoretically, α could not reach $\frac{3}{2}$ exactly in Sutherland's formula (10), so we pick this value as the upper limit. Thus, the Chapman's model and Sutherland's model of viscosity origin can be unified by Eq. (11) and extended at cosmic large scale also for late-stage evolutional era, while variations are provided mainly by the parameter α.

D. The red-shift related CMB temperature

The telling-tale CMB temperature, which is feasible to be measured at high red-shift in distant molecular clouds via absorption spectra [5], has provided us with lots of useful information on the universe evolution history. According to the current theories and observations, we are confident in assuming the linear relation between CMB temperature and its corresponding red-shift [6] holds:

$$T_{CMB}(z) \approx T_0(1 + z) \tag{12}$$

where $T_0 \approx 2.73$ the ambient CMB temperature measured at present. As we have discussed above, it is reasonable to combine Eq. (12) with the viscosity formulae (9, 10) in order to extend those concepts of viscosity from generally classical physics to cosmological physics.

II. MODELING AND CONSTRAINING

A. unified fluid model

1. *basic equations*

According to the discussion above, the general formula adopted as our temperature related viscosity model reads

$$\zeta = \zeta_0 T^{\alpha} \tag{13}$$

where ζ_0 is a simplified parameter and parameter α can be chosen as either $\frac{1}{2}$ (for Chapman's relation case) or $\frac{3}{2}$ (for Sutherland's formula limit case). In addition to the expression of viscosity, unified fluid model and linear relation between CMB temperature and the red-shift are considered together in this subsection. Now we can rewrite the mathematical expressions of those two premises as:

$$p = -3H\zeta \tag{14}$$

$$T(z) = T_0(1 + z) \tag{15}$$

By combining the two Einstein's FRW equations, i.e: Eqs. (1, 2) with our temperature related viscosity model: Eqs. (13)–(15) we have

$$\frac{3}{2}H^2+\dot{H}=12\pi GH\zeta_0T_0^{\alpha}(1+z)^{\alpha} \tag{16}$$

It is easy to convert the differential equation with respect to cosmic time into that with respect to red-shift z or scalar factor a, by utilizing

$$\dot{H}=-(1+z)H(\frac{\partial H}{\partial z})=H(\frac{\partial H}{\partial \ln a}) \tag{17}$$

The general solution reads

$$E(z)=\frac{e^{-\frac{\beta}{\alpha}(1+z)^{\alpha}}}{C-\frac{3}{2\alpha}Ei[-\frac{\beta}{\alpha}(1+z)^{\alpha}]} \tag{18}$$

where $\beta=\frac{12\pi G\zeta_0T_0^{\alpha}}{H_0}$ with H_0 represents the Hubble parameter at present. In order to nail down the integral constant C, we set the boundary or initial condition $E(z=0)=1$, so that $C=e^{-\frac{\beta}{\alpha}}+\frac{3}{2\alpha}Ei[-\frac{\beta}{\alpha}]$. The symbol Ei represents the exponential-integral-function $Ei(x)=-\int_x^{\infty}\frac{e^{-t}}{t}dt$. We briefly show the behavior curve of the $Ei(x)$ function in Fig. (1).



FIG. 1: The curve of function $Ei(x)$. A turning point occurs when $x=0$, and the left branch ($Ei(x<0)$) converges to zero as x goes to minus infinity. Since $-\frac{\beta}{\alpha}(1+z)^{\alpha}$ is always negative if we ask the redshift not less than -1, the value of Ei in Eq. (18) is confined under zero mainly.

Since the lower limit of viscosity is zero (viscosity free) or a constant (cosmological constant-like case), parameter β should be positive and so is the value of $-\frac{3}{2\alpha}Ei[-\frac{\beta}{\alpha}(1+z)^{\alpha}]$ when we consider the recent evolution of the universe, say from z=0.4 to the future. It is possible that Eq. (18) may suggest a mathematical and physical turning detour at the point $C=\frac{3}{2\alpha}Ei[-\frac{\beta}{\alpha}(1+z)^{\alpha}]$, and this case only happens when the value of C is negative (also see Fig. (1)) according to the discussion above. So the negative C may denote bouncing universe models while positive C indicates models without the bouncing solution character. If C equals to zero, α will be related to β by the boundary condition, and consequently with a strict control on the freedom of parameter β to make the model less arbitrary.

2. observational data constraints

To confront the models for extreme cases with the observational data-sets we use, for convenience, the reduced dimensionless Hubble parameter $h=H_0/(100\ kms^{-1}Mpc^{-1})$ at present with the parameter β. Since the matter-energy densities of dust matter and dark energy are coupled together in the unified fluid model, the observational H(z) and SNe Ia data sets (will be briefly discussed in the appendix) could be expected to give appropriate constraints on our model.

When the model with case $\alpha=\frac{1}{2}$, the observational H(z) and SNe Ia data sets together generate an overlap region with 1σ and 2σ confidence ranges for the parameter pair (β,h) (results are shown in the left panel of Figs. (2)).



FIG. 2: The 1 and 2 σ confidence ranges of (β, h), constrained by the observational H(z) and SNe Ia data sets. The left panel shows an ideal overlapped area for model with the $\alpha = 1/2$ case, while to constraints for the extreme model case $\alpha = 3/2$ the data-sets fittings from observational H(z) do not overlap with those from the union 2.1 of SNe Ia data-sets.

While at case $\alpha = \frac{3}{2}$, the H(z) and SNe Ia data-sets do not give overlap region for the parameter pair (β, h), (results are shown in the right one of Figs. (2)). Recall that when we first mentioned the case with $\alpha = \frac{3}{2}$, we take it as an extreme limit of the model parameter α in Sutherland's formula. Now we see that the limit could not only be reached theoretically, but also unfavored by observational data fittings.

We find out that the unified fluid models are with interesting features as the dark energy phenomena required. When the parameter $\alpha < 3/2$, the unified fluid model and the theoretical explanation of viscosity origin (the molecular-like interaction theory) are acceptable by the data fittings as shown. We keep in mind that so far by the global fitting any feasible cosmology model to include the dominated dark energy should not deviate the ΛCDM model away obviously. That conclusion leads us to move on to the study of effective viscosity functioning on the basis of the well-known ΛCDM model.

B. ΛCDM-V model

1. *basic equations*

In this subsection, we focus on the application of viscosity to the ΛCDM model. The expression of viscosity still follows Eq. (11) which contains both the physical content of Chapman's relation and Sutherland's formula, while the corresponding Einstein's component equations which include the constant dark energy Λ term in the flat FRW metric read

$$H^2 = \frac{8\pi G}{3}\rho = \frac{8\pi G}{3}(\rho_m + \rho_\Lambda) \tag{19}$$

$$H^2 + \dot{H} = -\frac{4\pi G}{3}(\rho + 3\frac{p}{c^2}) \tag{20}$$

where ρ_m is energy density of pressureless matter, ρ_Λ represents energy density of the constant dark energy, and c is the light speed in the vacuum (in natural units c equals to one). The expression of the effective pressure p including the viscosity contribution in Eq. (20) then reads

$$p = -\rho_\Lambda c^2 - 3H\zeta \tag{21}$$

where the first term is the effect of constant dark energy which EOS parameter $w_{DE} = -1$ exactly, and the second term comes from the effect of viscosity. We do not regard viscosity as one peculiar component of cosmic matter-energy in the universe, but as a gross grain approximation effect of the thermal and dynamical effect of interactions in the fluid which permeates the universe.

The combined form of Eqs. (19)–(21) reads

$$\dot{H} = -\frac{3}{2}\Omega_{m0} H_0^2 (1+z)^3 + 12\pi G H \zeta \tag{22}$$

which can be parameterized as

$$\frac{\partial E}{\partial z} = A\frac{(1+z)^2}{E} - \frac{B}{H_0}(1+z)^{\pm\frac{1}{2}} \tag{23}$$

where $A = \frac{3}{2}\Omega_{m0}$ and $B = 12\pi G \zeta_0 T_0^{\alpha}$ with α equals to $\frac{1}{2}$ or $\frac{3}{2}$ according to the specific formulas of viscosity. The power index $\pm\frac{1}{2}$ in the above equation corresponds to Sutherland's formula and Chapman's relation respectively. We name this model ΛCDM-V where V is short for the viscosity. For convenience, letter "a" and "b" are added as postfixes when α equals to $\frac{1}{2}$ and $\frac{3}{2}$ respectively. The new model differs from the standard ΛCDM with an additional term $\frac{B}{H_0}(1+z)^{\pm\frac{1}{2}}$ in Eq. (23), which has no clearly analytical solution. We will use numerical method by data statistic analysis in model constraining. Notice here exist two theoretical restrictions for Eq. (23), i.e: $E(z=0) = 1$ which is a natural boundary condition, and $B > 0$ which must be ensured due to the physical existence of viscosity.

2. observational data constraints

In this subsection we have proposed a model which explicitly announces that the universe consists of baryons, cold dark matter, and constant dark energy with viscosity, and both the dark energy and viscosity contributes to the effective pressure. In addition to the observational H(z) and SNe Ia data sets, the baryon acoustic oscillation (BAO) data set which has strong constraint on dark energy models will also be included in model constraining. (Applications of BAO data will be briefly discussed in the appendix and we should mention that several observation H(z) data points in the general H(z) data catalogue are from BAO. We will not treat them twice in doing the statistic analysis [18] for the models.)

The best-fit results for model Va and Vb are listed in Table (I), besides which the confidence ranges of parameter pair (Ω_{m0}, B) of both models are shown in Figs. (3).

TABLE I: Best-fit parameters of ΛCDM-V models by various data sets

data sets applied	H(z)	SNe	H(z)+BAO
model-Va			
χ^2_{min}	15.058	562.225	16.621
Ω_{m0}	0.367	0.283	0.284
$B(=12\pi G\zeta_0 T_0^{\frac{1}{2}})$	31.422	0.870	4.533
$H_0\ \ (km/s/Mpc)$	73.332	69.884	70.549
model-Vb			
χ^2_{min}	15.037	562.224	16.740
Ω_{m0}	0.589	0.290	0.288
$B(=12\pi G\zeta_0 T_0^{\frac{3}{2}})$	58.902	1.610	1.921
$H_0\ \ (km/s/Mpc)$	73.748	69.885	69.882

The observational H(z) or SNe Ia data alone does not provide much tight constraints on parameter pair (Ω_{m0}, B), until BAO data is taken into consideration. The overlapped 2σ confidence ranges in Figs. (3) favour positive value of B, and confine it under the order of the magnitude of 10, while the overlapped confidence ranges of Ω_{m0} are also acceptable, which lie between 2.5 and 3.0. The confidence ranges of H_0 are not shown here, since the constraints on the Hubble parameter by each individual data set are tight and give out similar values around 70 $km/s/Mpc$. Besides, among the three parameters, the values of B and Ω_{m0} matter most to our newly viscous cosmology model constraining.



FIG. 3: The 1σ and 2σ confidence ranges for parameter pair (Ω_{m0}, B), constrained by the observational H(z) (dashed line), SNe Ia (solid line) and BAO+H(z) data-sets (thick solid line). The contours of confidence ranges given by the observational H(z) and SNe Ia data-sets exceed the figures' boundary, but we are mainly interested in the overlapped area. Constraints from the observational H(z) data alone deviate a little from those of the other two, which indicates that the observational H(z) data lacks enough precision to be an independent data set (unlike the good quality of SNe Ia data-sets as the Union 2.1) for model constraining. But considering the future astrophysics observations progress the observational H(z) data is still our first choice when a quick fitting or an auxiliary data set is required.

III. DISCUSSIONS ON THE ΛCDM-V MODEL

In the SI units, the expression of viscosity reads clearly as

$$\zeta = \frac{Bc^2}{12\pi G}(1+z)^{\alpha} \tag{24}$$

where α equals 1/2 in model-Va and 3/2 for extreme limit in model-Vb, and $c = 2.99 \times 10^8\ m/s$ is the light speed in vacuum. The above equation can be expressed explicitly as

$$\zeta \approx 1.151 \times B(1+z)^{\alpha} \times 10^6 \quad Pa \cdot s \tag{25}$$

The evolutional behavior of viscosity is related to the expansion scalar $a = \frac{1}{1+z}$ which has been pointed out in Eq. (25). The magnitude of viscosity will be gradually reduced by the on-going expansion. This provide us an physically acceptable transition mechanism of fluid from imperfectness to perfectness during the cosmic evolution.

According to the results given by model constraining, both model Va and Vb are acceptable. The results provide an allowed range for the value of B, and consequently set an upper limit for viscosity parameter. Since in this present study we are mainly interested in the evolutional effect of viscosity on late stage cosmological evolution of the universe, in the following we will show how much deviation the viscosity model constrained by the background data-sets may provide against the standard scenario, the ΛCDM model.

For convenience, we re-express the ΛCDM model in a simple form with the Ω_{m0} as the cosmic observational matter composite fraction with repeat to the critical density of the observable universe today

$$H^2_{\Lambda CDM} = H_0^2[\Omega_{m0}(1+z)^3 + (1-\Omega_{m0})] \tag{26}$$

According to the latest WMAP-9yr observational results [1], we set the parameters $\Omega_{m0} = 0.287$ and $H_0 = 69.32$ by the global best-fit results for the standard cosmology model. In order to observe the effects of viscosity, we also set the parameter $A = 0.427$ and $H_0 = 70$ for model Va and Vb, while the value of parameter B varies from 0 to 10 roughly according to the previous fitting results.

The contrasts with H-z relation between the viscous models and standard cosmology models are illustrated in Figs. (4). Viscosity contribution suppresses the value of Hubble parameter at high red-shift, which means that it increases the cosmic age. As we know today that the age of the universe is 13.772 ± 0.059 Gyr as predicted by the standard cosmology model with observations [1]. For the possible largest value of parameter B, the standard cosmic age could be extended by 1.012 and 1.041 times by Va and Vb models respectively. Generally speaking, the age of the universe predicted by the ΛCDM-V models will not be older than an upper limit as 15 Gyr.

We are also interested in the fate of the universe predicted by the viscous models. To some extent, the ΛCDM-V model can be regard as a phantom-like dark energy model according to Eq. (21), (also see Figs. (5)). As the expansion goes on the observational universe will tend to infinities, the cosmic media viscosity will fade out or be negligent totally and the imperfect fluid will gradually become perfect one to satisfying accuracy.

There are basically four types of future singularity [7]. Type I (Big Rip): the scale factor becomes infinite at a finite time in the evolution future. Type II (Sudden Rip): the scale factor and energy remain finite at the rip time when pressure becomes infinite. Type III: the scale factor remains finite when energy and pressure become infinite in the finite future. Type IV: the scale factor remains finite in the finite future when energy and pressure vanish, and the higher derivatives of H diverge. We may extend the classfication to include the EoS singularity as well.

When the red-shift approaches -1, the Hubble parameter remains finite as shown in Figs. (4), which means the Big Rip is avoided by model Va and Vb respectively; according to Eqs. (21), (13) and (15), pressure will always be finite, so Type II and III singularities are also absent. The last kind of singularity is apparently not predicted, so the ΛCDM-V model will not experience the above four types of singularity as well as the EoS singularity obviously.

Taking into account of the no-rip, little rip and pseudo-rip [8] cases, there are seven types of fate for the universe evolution. The ΛCDM model certainly predicts the no-rip future. We have already known that during the expansion of the universe, our viscous model will gradually evolve into the standard perfect one. With the Hubble parameter predicted to be constant (dominated by a cosmology constant like term) when time goes to infinity, the ΛCDM-V models are free of the kind of rip destiny.



FIG. 4: The H-z relation diagram with H(z) data as reference background. The thick line represents the H(z) behavior from the ΛCDM model, while the shadowed area represents the possible H(z) values of the specific model with the value of B varying from range 0 to 10 for model comparings.

IV. CONCLUSIONS

In the above sections we have explored the possible effects of the existence of temperature related viscosity on cosmological evolution by proper modeling which is supported by observed data sets fittings. Similar research works in literature have proposed a variety of kinds of viscosity forms, from constant viscosity parameter [9] [11] to complicated turbulence phenomena related forms as possibly appearing in the complex cosmos media physics [12], and the temperature related viscosity in this present work is proved to be one possibility of them with suitable models supported by observational data sets well confronted.

In this article, at first we have attempted to conceive the unified dark energy (matter, dark matter and dark energy unified) models, expected that the temperature related viscosity may be one of the better candidates for dark energy phenomena. The observed data-sets fitting results have shown some inconsistence between the data-sets fitting from the SNe Ia confidence ranges against those from the observational H(z) data-sets for the viscous dark energy model with the extreme case of $\alpha = 3/2$. The not so precise observational H(z) data-sets fitting results (for this point we can see clearly also from the listed table one for the H(z) fitting results accordingly) for the specific model with parameter $\alpha = 3/2$ case (actually in the Sutherland's relation, the parameter $\alpha < 3/2$) imply we need further proper dark energy



FIG. 5: The w-w' phase space of the ΛCDM-V models, where the prime represents $\frac{\partial}{\partial z}$ instead of the $\frac{\partial}{\partial N}$ with N=lna(t) and w here is the effective EOS parameter since we hope to give a clear behavior of the effective equation of state especially when $w = -1$ rather than to illustrate the dark energy model classification by Caldwell and Linder parametrizasion (Refs. [13]). The thick line represents the w-w' curve of the ΛCDM model, while the dashed line shows the limit of viscous models in phantom area with value $B = 10$.

model building, which means this kind of viscosity dark energy could only be one possible component describing some properties for currently cosmic dark energy phenomena, while the model with $\alpha = 1/2$ case is satisfying. Similar researches have pointed to basically the same results, which indicate that unified fluid models with viscosity can mimic the evolutional behavior, but usually cause damping of density perturbations [9] [10].

Following the extension of our first trial we have then proposed the ΛCDM-V model, which is a modification of the standard ΛCDM model accordingly, turning the idealized perfect fluid cosmology model into the practically imperfect one with proper viscous energy-momentum tensor functioning and is fitted satisfactorily with the currently main observational data-sets. The upper limit on parameter B given by data-sets constraints is still loose, since when B equals to 10 the DE density may become negative at high red-shift, but notice that the null energy condition (NED) is not violated. That is mainly caused by the data-sets whose observational range are limited under red-shift $z = 2$. Since late-stage evolution was mainly discussed, CMB-shift parameter has not been taken into account, which can provide a richer content about the past history of the universe. For consistency, we strongly suggest that a theoretically proper upper limit on B may lie around the magnitude of order of 0.1 in order to avoid that crossing phenomenon of DE energy density. Introducing interaction between DE and DM into this model may also expected to raise the upper limit. For illustration, the evolution behavior of DE density of ΛCDM-V model with $B = 0.1$ are shown in Fig. (6).



FIG. 6: The density evolution of DE, when $B = 0.1$ as an example, without singularity or crossing behavior at least after the occurrence of decoupling ($z < 1100$).

In the Figs. (5) for numerical comparisons, we can see the parameterized equation of state (EoS) parameter evolution lines of the ΛCDM-V and the standard ΛCDM model can match at present stage and in the evolution future, to form a consistent picture which is in accordance with the no-rip future predicted by the new model. In addition to increasing the cosmic age and being free of future singularity, we would like to say the smooth transition mechanism is another pleasing feature of the ΛCDM-V model. In previous researches conducted in some cosmological dark energy model buildings, imperfect fluid models remain to be with imperfect fluid all along the evolution of the universe. In this present work, the model Va and Vb could converge to the cosmic perfect fluid model, the globally fitted well ΛCDM model, as the cosmic expansion goes on to infinity to dilute the viscosity effects, by providing a smooth transition mechanism from practically imperfect fluid models to perfect fluid ones.

Considering about the complexities and difficulties for the physical viscous media descriptions in observational universe, in the present work we have not included the density perturbation discussions which may give much tighter constraints on model buildings, like via the integrated Sachs-Wolfe effect or matter power spectrum simulations (see Refs. [9] [11] [14] [15]). We strongly believe that more precise observed data sets given by future astrophysics observation instrumental achievements can certain provide higher capacity performance in model constraining and constructing and will finally pin down the physics for long puzzling cosmic dark matter problems and mysterious dark energy phenomena.

The currently possible origins and explanations of viscosity physics may remind us of that trying to express viscosity effect in only one form could be too simple. The rich viscosity effects in cosmos may be an assemblage of a variety of dynamical and kinematics effects of locally cosmic motion and globally cosmos large-scale evolution, which may show different features when observed on different scales and /or in different evolutional stages. We may say this present article research reveals only a tip of the whole iceberg. Nevertheless, continuous study endeavors to the mysterious dark sector physics will surely shed light towards the fundamental understandings of our universe.

Acknowledgement

Interesting communications with Profs. S. Odintsov and Lewis H. Ryder are very enjoyable. We thank Prof. J. D. Barrow for kindly pointing out Refs. [10] to us and Dr. Hermano Velten for offering precious suggestions. We also appreciate programming helps from Xiaolong Du, and beneficial discussions with Han Dong.

This work is partly supported by Natural Science Foundation of China under Grant Nos.11075078 and 10675062, and by the project of knowledge Innovation Program (PKIP) of Chinese Academy of Sciences (CAS) under the grant No. KJCX2.YW.W10 through the KITPC astrophysics and cosmology program where we have initiated this present work.

Appendix

For the possible benefits to the concerned academic community with our hope we write out in details the data fitting preparations in this work. We employ heavily in this presentation the χ^2 statistical method [16] [17] [18] for model buildings with probability distribution function which reads (for measuring confidence range of parameters or coefficients)

$$P(\vartheta) = \mathcal{N} e^{-\frac{\chi^2(\vartheta)}{2}} \tag{27}$$

where $\mathcal{N}$ is the normalization coefficient to ensure $\int P(\vartheta) d\vartheta = 1$.

A. Observed Hubble Parameter data (OHD) sets (the observational H(z) data sets)

The H(z) data-sets contains 28 measurement points so far. It provides a clear and direct picture of $H(z)$ evolution over the range $0.1 < z < 1.5$. OHD depicts the relation between the Hubble parameter H(z) and the according red-shift z, which is also related and described directly when conceiving cosmic models. The value catalogue of this data-set are reported in Table (II). The χ^2 calculation for OHD reads

$$\chi^2_{OHD} = \sum_{i=1}^{28} [\frac{H_{obs}(z_i) - H_0 E_{th}(z_i;\vartheta)}{\sigma_i}]^2 \tag{28}$$

TABLE II: Observed Hubble parameter Data, Refs. [19–24]

z	H (km/s/Mpc)	error
0.090	69	12
0.170	83	8
0.270	77	14
0.400	95	17
0.900	117	23
1.300	168	17
1.430	177	18
1.530	140	14
1.750	202	40
0.480	97	62
0.880	90	40
0.179	75	4
0.199	75	5
0.352	83	14
0.593	104	13
0.680	92	8
0.781	105	12
0.875	125	17
1.037	154	20
0.24	79.69	3.32
0.43	86.45	3.27
0.07	69.0	19.6
0.12	68.6	26.2
0.20	72.9	29.6
0.28	88.8	36.6
0.44	82.6	7.8
0.60	87.9	6.1
0.73	97.3	7.0

B. Type Ia Supernova (SNe) data set (Union 2.1)

SNe Ia data is believed to provide a solid piece of evidence for cosmic acceleration at present era of evolution of the Universe. Type Ia supernova are regarded as standard candles for distance measurement in observation universe, with same absolute magnitude $M \approx -19.3 \pm 0.3$. Union2.1 data set which is used in this paper contains 580 measurement points. The theoretical distance modulus is defined as

$$\mu_{th} = m_{th} - M = 5log_{10}D_L(z) + \mu_0 \tag{29}$$

where $D_L = (1+z)\int_0^z \frac{dz'}{E(z')}$, under the assumption that our universe is spatially flat, at least within the space-time we are able to have observed, $\Omega_k = 0$; and $\mu_0 = 42.38 - 5log_{10}h$,, where h is the dimensionless reduced Hubble parameter at present. The χ^2 fitting formula for SNe Ia data then reads:

$$\chi^2_{SNe} = \sum_{i=1}^{580}[\frac{\mu_{obs}(z_i) - \mu_{th}(z_i;\vartheta)}{\sigma_i}]^2 \tag{30}$$

Since some of our viscous models are very complicated, we may reformulate the SNe Ia data set by converting the $\mu \sim z$ relation into $D_L \sim z$ relation through the following equations:

$$D_L(z_i) = 10^{\frac{\mu(z_i)-\mu_0}{5}}, \tag{31}$$

besides, we should convert the old error bar value of each measurement point into the corresponding new error value with

$$\sigma_{new}(i) = D_L(z_i)[10^{\frac{\sigma_{old}(i)}{5}} - 1]. \tag{32}$$

Under this transformation of data sets, we are free from the Log_{10} calculation while experimenting χ^2 fitting. Thus the reformed χ^2 calculation is

$$\chi^2_{SNe} = \sum_{i=1}^{580}[\frac{D_{Lobs}(z_i) - D_{Lth}(z_i;\vartheta)}{\sigma_{new}(i)}]^2 \tag{33}$$

C. Baryon Acoustic Oscillations (BAO) data set

The BAO is originated from perturbated sound waves in the early universe when primordial sound wave could travel at relativistic speed through space-time. We have gathered data from 6dFGS [25] (Beutler *et al.* 2011), Wiggle Z [26] (Blake *et al.*2011), and SDSS [27] (Eisenstein *et al.*2005). The data sets will be listed in Table (III) below. The statistically χ^2 analysis for BAO data-sets reads

$$\chi^2_{BAO} = \sum_{z_i}[\frac{A_{obs}(z_i) - A_{th}(z_i;\vartheta)}{\sigma_A}]^2, \tag{34}$$

where

$$A(z) = \sqrt{\Omega_{m0}}E_z^{-1/3}[\frac{1}{z}\int_0^z \frac{dz'}{E(z')}]^{2/3} \tag{35}$$

TABLE III: Baryon Acoustic Oscillation Data, Refs. [25–27]

z	A	error
0.106	0.526	0.028
0.35	0.469	0.017
0.6	0.452	0.018

---

[1] G. Hinshaw *et al.*, arXiv:1212.5226 (2012).
[2] Gong-Bo Zhao, *et al.*, Phys. Rev. Lett. **109**, 171301 (2012).
[3] Xin-He Meng, Zhi-Yuan Ma, Eur. Phys. J. C **72**, 2053 (2012).
Xin-He Meng, Jie Ren, Ming-Guang Hu, Commun. Theor. Phys. **47**, 379-384 (2007).
Xu Dou, Xin-He Meng, Adv. Astron. **2011**, 829340 (2011).
[4] T. Buchert, S. Rasanen, Ann. Rev. Nucl. Part. Sci. **62**, 57-59 (2012).
Nan Li *et al.*, Fortsch. Phys. **56**, 465-474 (2008).
[5] P. Noterdaeme, *et al.*, arXiv:1012.3164.
[6] J. M. LoSecco, G. J. Mathews, Yun Wang, Phys. Rev. D **64**, 123002 (2001).
[7] Shin'ichi Nojiri, *et al.*, Phys. Rev. D **71**, 063004 (2005).
[8] P. H. Frampton, *et al.*, Phys. Rev. D **85**, 083001 (2012).
[9] H. Velten, D. J. Schwarz, JCAP. **1109**, 016 (2011), arXiv:1107.1143v2.
[10] B. Li, J. D. Barrow, Phys. Rev. D **79** 103521 (2009).
J. D. Barrow, Phys. Lett. B **180**, 335 (1986).
J. D. Barrow, Nucl. Phys. B **310**, 743 (1988).
J. D. Barrow, String-driven Inflation, in *The Formation and Evolution of Cosmic Strings*, eds. G. Gibbons, S. W. Hawking, T. Vachaspati, pp 449-464, CUP, Cambridge (1990).
[11] H. Velten, D. J. Schwarz, Phys. Rev. D **86**, 083501 (2012) arXiv:1206.0986v3.
[12] I. Brevik, *et al.*, Phys. Rev. D **86**, 063007 (2012).
[13] R. R. Caldwell, E. V. Linder, Phys. Rev. Lett. **95**, 141301 (2005).
R. R. Caldwell, Phys. Lett. B **545** 23-29 (2002).
Y. D. Xu *et al.*, Astrophys. Space. Sci. **337**, 493-498 (2012).
[14] J. C. Fabris, H. E. S. Velten, W. Zimdahl, Phys. Rev. D **81**, 087303 (2010), arXiv:1001.4101v1.
[15] J. C. Fabris, H. E. S. Velten, Phys. Lett. B **694**, 289-293 (2011), arXiv:1007.1011v1.
[16] Hao Wei, JCAP. **1104**, 022 (2011), arXiv:1012.0883v3.
[17] S. Capozziello,S. Nojiri, S. D. Odintsov, *et al.*, Phys. Rev. D **73**, 043612 (2006) arXiv:astro-ph/0508350v3 (2006).
[18] Meng Xin-He, Du Xiao-Long, Commun. Theor. Phys. **57**, 227-233 (2012)
Xinhe Meng, Xiao-Long Du, Phys. Lett. B **710**, 493-499 (2012)
[19] J. Simon, L. Verde, R. Jimenez, Phys. Rev. D **71**, 123001 (2005).
[20] D. Stern, R. Jimenez, L. Verde, M. Kamionkowski, S. A. Standford, J. Cosmology Astropart. Phys, **02**, 008 (2010).
[21] M. Moresco, A. Cimatti, R. Jimenez, *et al.* J. Cosmology Astropart. Phys., arXiv:1201.3609v1. (2012).
[22] E. Gaztanaga, A. Cabre, L. Hui, MNRAS, **399**, 1663 (2009).
[23] Cong Zhang, Han Zhang, Shuo Yuan, Tong-Jie Zhang, Yan-Chun Sun, arXiv:1207.4541 (2012).
[24] C. Blake,S. Brough,M. Colless *et al.*, Mon. Not. Roy. Astron. Soc., **425**, 405, arXiv:1204.3674 (2012).
[25] Florian Beutler *et al.*, Mon. Not. R. Astron. Soc. **416**, 3017-3032 (2011), arXiv:1106.3366v1.
[26] Chris Blake *et al.*, Mon. Not. R. Astron. Soc. **415**, 2892-2909 (2011), arXiv:1105.2862v1.
[27] Daniel J. Eisenstein *et al.*, Astrophys. J. **633**, 560-574 (2005), arXiv:astro-ph/0501171v1.
[28] Mi-Xiang Lan, Miao Li, *et al.*, Phys. Rev. D **82**, 023516 (2010) arXiv:1002.0978v3 (2010).
[29] D. H. Winberg *et al.*, arXiv:1201.2434v1 (2012).
[30] W. Hu and N. Sugiyama, Astrophys. J. **471**, 542 (1996).
[31] Andrew Liddle, *An Introduction to Modern Cosmology 2ed* (Weiley, 2003).
[32] S. Chapman, T. G. Cowling, *The Mathematical Theory of Non-uniform Gases 3rd* (Cambridge, 1995).
[33] L. D. Landau, E. M. Lifshitz, *Fluid Mechanics 2ed* (Elsevier, 2004).
[34] Luca Amendola, Shinji Tsujikawa, *Dark Energy: Theory and Observations* (Cambridge, 2010)